

## **Background:**

We are facing a biodiversity crisis that is unprecedented in human history, caused by myriad human impacts including habitat loss and fragmentation, invasive species, pollution, unsustainable harvesting, and climate change (Barnosky et al. 2011). For example, the 2018 World Wildlife Fund Living Planet Index (WWF 2018) suggests that we have lost approximately 70% of global diversity since 1970 – a period of marked human population growth and dramatic increases in industrialization and economic growth in many countries. The UN Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES) estimates that there are one million species at risk of extinction in the next decades (IPBES 2019). Conservation action (e.g. national and international treaties, species legal protections, reserve design and designation) requires detailed quantitative data on temporal trends in population sizes (e.g. Bonebrake et al. 2010), shifts in species ecology (e.g. breeding phenology; Adams 2010), and changes to species distributions (e.g. Sekercioglu et al. 2008; Marini et al. 2009). The need for continuous monitoring and demographic data is particularly important for migratory species, especially in the Northern Hemisphere, that require breeding grounds, stop-over sites during seasonal migrations, and wintering grounds that may encompass many countries (e.g. Amano et al. 2010; Wang et al. 2017). Birds also serve as important bioindicators of ecosystem health surveys of waterbird diversity in particular can provide insights into ecosystem health (Zhang and Ma 2011).

One of the most important migratory stop-over sites in eastern Asia is the Chongming Dongtan Nature Reserve, located on the eastern edge of Chongming Island, the largest estuarine alluvial island in the world and part of the Yangtze River Delta (MacKinnon et al. 2012). Reflecting its global ecological importance, the Dongtan Reserve is designated as National Nature Reserve in China and a Ramsar Wetland of International Importance (MacKinnon et al. 2012). Dongtan wetlands are critical for over 100 migratory waterbirds, including the critically endangered spoon-billed sandpiper (*Eurynorhynchus pygmeus*), the endangered black-faced spoonbill (*Platalea minor*) and the vulnerable hooded crane (*Grus monacha*) (MacKinnon et al. 2012).

There are many means to monitor population numbers for individual species or groups of species that includes visual monitoring by human observers, automated audio recorders with song recognition, and environmental DNA surveys. This study will be part of a larger collaborative project to compare different methods for surveying avian diversity and the environments upon which they depend including invertebrate diversity and water chemistry.

## **Objectives**

Most migratory species in stop-over sites like Dongtan Reserve forage on benthic invertebrates to build energy for further migration (Zhu et al. 2007). Benthic invertebrates also serve as an excellent indicator of water quality and ecosystem health (e.g. Azrina et al. 2006). The overarching goal is to quantify changes in invertebrate diversity and abundance over the course of spring bird migration at

Dongtan Reserve using sampling of the sediment and associated benthos, and subsequently using a series of taxonomic keys to identify taxa. The hypothesis is that maximum bird migrant abundance and species richness will coincide with peak invertebrate abundance and biomass.

### **Significance of the project**

Humans have never had greater impacts on global environments than now. This is a consequence of burgeoning human populations and increasing industrialization in emerging economies like those in many Asian countries resulting in unprecedented rates of habitat loss and conversion, pollution, and ultimately global climate change. This is having profound and potentially irrevocable consequences for the natural world. More broadly, and in conjunction with other members of the MITACS research team, this study will help to better monitor aquatic ecosystem health using a suite of tools moving us to more comprehensive methods for monitoring the Yangtze River and other major riverine and lake systems.

## Timeline

TBD

### Literature cited

- Adams RA (2010) Bat reproduction declines when conditions mimic climate change projections for western North America. *Ecology* 91: 2437-2445.
- Amano T, Székely T, Koyama K, Amano H, Sutherland WJ (2010) A framework for monitoring the status of populations: an example from wader populations in the east Asian-Australasian flyway. *Biol Conserv* 143: 2238-2247.
- Azrina MZ, Yap CK, Ismail AR, Ismail A, Tan SG (2006) Anthropogenic impacts on the distribution and biodiversity of benthic macroinvertebrates and water quality of the Langat River, Peninsular Malaysia. *Ecotoxicol Environ Saf* 64: 337-347.
- Barnosky AD, Matzke N, Tomiya S, Wogan GOU, Swartz B, Quental TB, et al. (2011) Has the Earth's sixth mass extinction already arrived? *Nature* 471: 51-57.
- Bonebrake TC, Christensen J, Boggs CL, Ehrlich PR (2010) Population decline assessment, historical baselines, and conservation. *Conserv Lett* 3: 371-378.
- IPBES (2019) Summary for policymakers of the global assessment report on biodiversity and ecosystem services of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. S. Díaz, J. Settele, E. S. Brondizio E.S., H. T. Ngo, M. Guèze, J. Agard, A. Arneth, P. Balvanera, K. A. Brauman, S. H. M. Butchart, K. M. A. Chan, L. A. Garibaldi, K. Ichii, J. Liu, S. M. Subramanian, G. F. Midgley, P. Miloslavich, Z. Molnár, D. Obura, A. Pfaff, S. Polasky, A. Purvis, J. Razzaque, B. Reyers, R. Roy Chowdhury, Y. J. Shin, I. J. Visseren-Hamakers, K. J. Willis, and C. N. Zayas (eds.). IPBES secretariat, Bonn, Germany.
- MacKinnon J, Verkuil YI, Murray N (2012) IUCN situation analysis on East and Southeast Asian intertidal habitats, with particular reference to the Yellow Sea (including the Bohai Sea). Occasional Paper of the IUCN Species Survival Commission No.47. IUCN, Gland, Switzerland and Cambridge, UK. ii + 70 pp.
- Marini MÂ, Barbet-Massin M, Lopes LE, Jiguet F (2009) Predicted climate-driven bird distribution changes and forecasted conservation conflicts in a neotropical savanna. *Conserv Biol* 25: 1558-1567.
- Sekercioglu CH, Schneider S, Fay JP, Loarie SR (2008) Climate change, elevational range shifts, and bird extinctions. *Conserv Biol* 22: 140-150.
- Wang W, Fraser JD, Chen J (2017) Wintering waterbirds in the middle and lower Yangtze River floodplain: changes in abundance and distribution. *Bird Conserv Int* 27: 167-186.
- WWF (2018) Living Planet Report - 2018: Aiming Higher. Grooten, M. and Almond, R.E.A.(Eds). WWF, Gland, Switzerland.

- Zhang, W, Ma J (2011) Waterbirds as bioindicators of wetland heavy metal pollution. *Proc Environ Sci* 10: 2769-2774
- Zhu J, Jing K, Gan X, Ma Z (2007) Food supply in intertidal area for shorebirds during stopover at Chongming Dongtan, China. *Acta Ecol Sinica* 27: 2149-2159.